

RELIATech GmbH
Lindenerstr. 15
38300 Wolfenbüttel
Germany

Tel.: +49 5331 8586 987
Fax.: +49 5331 8586 989
Email: info@reliatech.de
web: www.reliatech.de

Recombinant Human PDGF-AB

Description: PDGFs are disulfide-linked dimers consisting of two 12.0-13.5 kDa polypeptide chains, designated PDGF-A and PDGF-B chains. The three naturally occurring PDGFs; PDGF-AA, PDGF-BB and PDGF-AB, are potent mitogens for a variety of cell types including smooth muscle cells, connective tissue cells, bone and cartilage cells, and some blood cells. The PDGFs are stored in platelet α -granules and are released upon platelet activation. The PDGFs are involved in a number of biological processes, including hyperplasia, chemotaxis, embryonic neuron development, and respiratory tubule epithelial cell development. Two distinct signaling receptors used by PDGFs have been identified and named PDGFR- α and PDGFR- β . PDGFR- α is high-affinity receptor for each of the three PDGF forms. On the other hand, PDGFR- β interacts with only PDGF-BB and PDGF-AB. Recombinant human PDGF-AB is a 25.5 kDa disulfide-linked dimer, consisting of one A chain and one B chains (234 total amino acids).

Source:	E.coli
Molecular Weight:	25.5 kDa
Purity:	>95% by SDS-PAGE and visualised by silver stain
Endotoxin level:	< 0.1 ng per ug of PDGF-AB
Stabilizer:	none
Buffer:	50 mM acetic acid
Formulation:	lyophilized

Biological Activity: The ED50 as determined by the dose-dependent stimulation of thymidine uptake by BALB/c 3T3 cells is < 1 ng/ml, corresponding to a specific activity of > 1 x 10⁶ units/mg.

Stability: The lyophilized PDGF-AB is stable for a few weeks at room temperature, but best stored at -20°C. Reconstituted PDGF-AB is best stored at -20°C to -70°C. **Avoid repeated freeze-thaw cycles.**

Usage: Human PDGF-AB is offered for research use. Not for drug use. **Not for human use.**

Catalogue number: 200-053S	Size: 2 μ g
	Range: 0.2-10.0 ng/ml

****please note: always centrifuge product before opening vial ****