

RELIATech GmbH
Lindenerstr. 15
38300 Wolfenbüttel
Germany

Tel.: +49 5331 8586 987
Fax.: +49 5331 8586 989
Email: info@reliatech.de
web: www.reliatech.de

Recombinant Human GM-CSF

Description: Granulocyte Macrophage Colony Stimulating Factor (GM-CSF), a 14,6 kDa protein consisting of 127 amino acid residues (Ala18-Glu144) and fused to a C-terminal His-tag (6x His), is a potent species specific stimulator of bone marrow cells and several other cell types.

GM-CSF was initially characterized as a growth factor that can support the *in vitro* colony formation of granulocyte-macrophage progenitors. It is produced by a number of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cells and fibroblasts) in response to cytokine or immune and inflammatory stimuli. Besides granulocyte-macrophage progenitors, GM-CSF is also a growth factor for erythroid, megakaryocyte and eosinophil progenitors. On mature hematopoietic cells, GM-CSF is a survival factor for and activates the effector functions of granulocytes, monocytes/macrophages and eosinophils. GM-CSF has also been reported to have a functional role on non-hematopoietic cells. It can induce human endothelial cells to migrate and proliferate. Additionally, GM-CSF can also stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma and adenocarcinoma cell lines. GM-CSF is species specific and human GM-CSF has no biological effects on mouse cells. GM-CSF exerts its biological effects through binding to specific cell surface receptors. The high affinity receptors required for human GM-CSF signal transduction have been shown to be heterodimers consisting of a GM-CSF-specific α chain and a common β chain that is shared by the high-affinity receptors for IL-3 and IL-5.

Source:	Insect cells
Molecular Weight:	14.6 kDa
Purity:	> 98% by SDS-PAGE and visualized by Silver stain
Endotoxin level:	< 0.1 ng per μ g of GM-CSF
Stabilizer:	none
Buffer:	PBS
Formulation:	lyophilized

Biological Activity: The ED_{50} as determined by the dose-dependent stimulation of the proliferation of human TF-1 cells is ≤ 0.1 ng/ml.

Reconstitution: The lyophilized GM-CSF is soluble in water and most aqueous buffers. The lyophilized powder can be reconstituted in water to a concentration of 0.1 mg/ml. This solution can be diluted into other buffered solutions or stored at -20°C for future use. For most *in vitro* applications, human GM-CSF exerts its biological activity in the concentration range of 0.05 to 1.0 ng/ml.

Stability: The lyophilized powder though stable at room temperature, is best stored desiccated below 0°C . Reconstituted GM-CSF should be stored in working aliquots at -20°C . **Avoid repeated freeze-thaw cycles.**

Usage: Human GM-CSF is offered for research use. Not for drug use. **Not for human use.**

Catalogue number: 400-010

Size: 5 μ g

Range: 0.05-1.0 ng/ml

****please note: always centrifuge product before opening Vial****