

RELIA*Tech* GmbH
Lindenerstr. 15
38300 Wolfenbüttel
Germany

Tel.: +49 5331 8586 987
Fax.: +49 5331 8586 989
Email: info@reliatech.de
web: www.reliatech.de

Antigen-Affinity Purified Polyclonal Antibodies

Anti-human sCD40 Ligand

Description: Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hsCD40L (human sCD40L). Anti-hsCD40L specific antibody was purified by affinity chromatography employing immobilized hsCD40L matrix.

Host species:	Goat
Antigen:	Recombinant human sCD40 Ligand
Purification:	Affinity chromatography
Stabilizer:	none
Buffer:	PBS; pH 7.2
Formulation:	lyophilized

Reconstitution/Stability: The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile water to a concentration of ≥ 0.2 mg/ml the antibody is stable for at least six weeks at 2-4°C.

Applications

Western Blot: To detect hsCD40-Ligand by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 $\mu\text{g/ml}$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hsCD40-Ligand is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

ELISA: To detect hsCD40-Ligand by direct ELISA (using 100 μl /well antibody solution) a concentration of at least 0.5 $\mu\text{g/ml}$ of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2-0.4 ng/well of recombinant hsCD40-Ligand.

Usage: Anti human sCD40 Ligand is offered for research use. Not for drug use. **Not for human use.**

Catalogue number: 102-P108G

Size: 100 μg