

**RELIA**Tech GmbH  
Lindenerstr. 15  
38300 Wolfenbüttel  
Germany

Tel.: +49 5331 8586 987  
Fax.: +49 5331 8586 989  
Email: [info@reliatech.de](mailto:info@reliatech.de)  
web: [www.reliatech.de](http://www.reliatech.de)

## Antigen-Affinity Purified Polyclonal Antibodies

# Anti-human FGF-16

**Description:** Produced from sera of goats pre-immunized with highly pure (>98%) recombinant human FGF-16 (human Fibroblast Growth Factor-16). Anti-human FGF-16 specific antibody was purified by affinity chromatography employing immobilized human FGF-16 matrix.

<b>Host species:</b>	Goat
<b>Antigen:</b>	Recombinant human FGF-16
<b>Purification:</b>	Affinity chromatography
<b>Stabilizer:</b>	none
<b>Buffer:</b>	PBS; pH 7.2
<b>Formulation:</b>	lyophilized

**Reconstitution:** Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Stability:** The lyophilized antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. **Avoid repeated freeze-thaw cycles.**

### Applications

**Western Blot:** To detect human FGF-16 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant human FGF-16 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

**ELISA: Indirect:** To detect human FGF-16 by indirect ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant human FGF-16.

**Sandwich:** To detect human FGF-16 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant human FGF-16.

**Usage:** Anti human FGF-16 is offered for research use. Not for drug use. **Not for human use.**

**Catalogue number:** 102-P125G

**Size:** 100 µg

**\*\*Please note: always centrifuge product before opening vial.\*\***