

RELIATech GmbH
Lindenerstr. 15
38300 Wolfenbüttel
Germany

Tel.: +49 5331 8586 987
Fax.: +49 5331 8586 989
Email: info@reliatech.de
web: www.reliatech.de

Antigen-Affinity Purified Polyclonal Antibodies

Anti-rat IFN-gamma

Description: Produced from sera of goats pre-immunized with highly pure (>98%) recombinant Rat IFN- γ . Anti-Rat IFN- γ specific antibody was purified by affinity chromatography employing immobilized Rat IFN- γ matrix.

Host species:	Goat
Antigen:	Recombinant rat IFN- γ
Purification:	Affinity chromatography
Stabilizer:	none
Buffer:	PBS; pH 7.2
Formulation:	lyophilized

Reconstitution: Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Stability: The lyophilized antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. **Avoid repeated freeze-thaw cycles.**

Applications

Neutralization: To yield one-half maximal inhibition [ND_{50}] of the biological activity of Rat IFN- γ (0.3 ng/ml), a concentration of 0.006-0.01 μ g/ml of this antibody is required.

ELISA: Indirect: To detect Rat IFN- γ by indirect ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IFN- γ .

Sandwich: To detect Rat IFN- γ by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IFN- γ .

Western Blot: To detect Rat IFN- γ by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat IFN- γ is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Usage: Anti rat IFN- γ is offered for research use. Not for drug use. **Not for human use.**

Catalogue number: 104-P03G

Size:

100 μ g

**** please note: always centrifuge vials before opening ****