



# Recombinant Human TGF-beta2

20190131BB

**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

<b>Cat.-no.:</b>	<b>100-105</b>
Size:	10 µg
Lot. No.:	According to product label

## Scientific Background

<b>Gene-ID (NCBI):</b>	7042
<b>Synonyms:</b>	TGFB2; TGF-beta2

The three mammalian isoforms of TGF- $\beta$ , TGF- $\beta$ 1,  $\beta$ 2,  $\beta$ 3, signal through the same receptor and elicit similar biological responses. They are multifunctional cytokines that regulate cell proliferation, growth, differentiation and motility as well as synthesis and deposition of the extracellular matrix. They are involved in various physiological processes including embryogenesis, tissue remodeling and wound healing. They are secreted predominantly as latent complexes which are stored at the cell surface and in the extracellular matrix. The release of biologically active TGF- $\beta$  isoform from a latent complex involves proteolytic processing of the complex and /or induction of conformational changes by proteins such as thrombospondin-1. TGF- $\beta$ 2 has been shown to exert suppressive effects on IL-2 dependent T-cell growth, and may also have an autocrine function in enhancing tumor growth by suppressing immuno-surveillance of tumor development. Recombinant human TGF- $\beta$ 2 is a 25.0 kDa protein composed of two identical 112 amino acid polypeptide chains linked by a single disulfide bond.

## Sequence

```
ALDAAYCFRN VQDNCCLRPL YIDFKRDLGW KWIHEPKGYN  
ANFCAGACPY LWSSDTQHSR VLSLYNTINP EASASPCCVS  
QDLEPLTILY YIGKTPKIEQ LSNMIVKSKC CS
```

## Database References

<b>Protein RefSeq:</b>	NP_001129071
<b>Uniprot ID:</b>	P61812
<b>mRNA RefSeq:</b>	NM_001135599

## Product Specifications

<b>Expressed in</b>	Insect cells
<b>Purity</b>	> 95% by SDS-PAGE & HPLC analyses
<b>Structural Information</b>	homodimer
<b>Endotoxin level</b>	< 0.1 ng /µg of protein (<1EU/µg).
<b>Formulation</b>	lyophilized
<b>Length (aa):</b>	112
<b>MW:</b>	25 kDa

**Stability:** The lyophilized protein is stable at room temperature for 1 month and at 4°C for 6 months. Reconstituted working aliquots are stable for 1 week at 2°C to 8°C and for 10 months at -20°C to -80°C.

**Reconstitution:** Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. *Do not vortex.* This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

**AVOID REPEATED FREEZE AND THAW CYCLES!**

**Biological Activity:** Determined by its ability to inhibit the mouse IL-4-dependent proliferation of mouse HT-2 cells. The ED<sub>50</sub> was found to be  $\leq 0.2$  ng/ml ( $\geq 5 \times 10^6$  units/mg).