



Recombinant Mouse IGF-1

20180712BB



FOR RESEARCH ONLY! NOT FOR HUMAN USE!

Cat.-no.:	M10-011
Size:	50 µg
Lot. No.:	According to product label

Sequence

GPETLCGAEL VDALQFVCGP RGFYFNKPTG YGSSIRRAPO
TGIVDECCFR SCDLRRLEMY CAPLKPTKAA

Database References

Protein RefSeq:	NP_034642.2
Uniprot ID:	P05017
mRNA RefSeq:	NM_010512

Scientific Background

Gene-ID (NCBI):	16000
Synonyms:	Igf1; Igf-1; Igf-I; C730016P09Rik

The IGFs are mitogenic polypeptide growth factors that stimulate the proliferation and survival of various cell types including muscle, bone, and cartilage tissue in vitro. IGFs are predominantly produced by the liver, although a variety of tissues produce the IGFs at distinctive times. The IGFs belong to the Insulin gene family, which also contains insulin and relaxin. The IGFs are similar by structure and function to insulin, but have a much higher growth-promoting activity than insulin. IGF-II expression is influenced by placenta lactogen, while IGF-I expression is regulated by growth hormone. Both IGF-I and IGF-II signal through the tyrosine kinase type I receptor (IGF-IR), but, IGF-II can also signal through the IGF-II/Mannose-6-phosphate receptor. Mature IGFs are generated by proteolytic processing of inactive precursor proteins, which contain N-terminal and C-terminal propeptide regions. Recombinant murine IGF-I is a 7.6 kDa globular protein containing amino acids including 3 intra-molecular disulfide bonds.

Product Specifications

Expressed in	E. coli
Purity	> 98% by SDS-PAGE & HPLC analyses
Endotoxin level	< 0.1 ng /µg of protein (<1EU/µg).
Formulation	lyophilized
Length (aa):	70
MW:	7.6 kDa

Stability: The lyophilized protein is stable at room temperature for 1 month and at 4°C for 6 months. Reconstituted working aliquots are stable for 1 week at 2°C to 8°C and for 12 months at -20°C to -80°C.

Reconstitution: Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. *Do not vortex.* This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.



AVOID REPEATED FREEZE AND THAW CYCLES!

Biological Activity: The ED₅₀ as determined by a cell proliferation assay using FDC-P1 cells is ≤ 2.0 ng/ml, corresponding to a specific activity of ≥ 5 x 10⁵ units/mg.