



Anti-human VEGFR-2/KDR-FITC (#EIC)

20191127DS



FOR RESEARCH ONLY! NOT FOR HUMAN USE!

Cat.-no.:	101-M20S-FITC
Size:	25 µg (in 50 µl)
Concentration:	500 µg/ml
Lot. No.:	According to product label
Country of origin:	Germany

Preparation: Monoclonal antibodies were produced with the help of BALB/c mice using recombinant human soluble extracellular KDR (D7) as the immunizing antigen

Target Background

Synonyms:	Vascular endothelial growth factor receptor 2
------------------	---

Disruption of the precise balance of positive and negative molecular regulators of blood and lymphatic vessel growth can lead to myriad diseases. Although dozens of natural inhibitors of hemangiogenesis have been identified, an endogenous selective inhibitor of lymphatic vessel growth has not to our knowledge been previously described. A splice variant of the gene encoding vascular endothelial growth factor receptor-2 (VEGFR-2) that encodes a secreted form of the protein, designated endogenous soluble VEGFR-2 (esVEGFR-2/KDR) has been described. The endogenous soluble esKDR inhibits developmental and reparative lymphangiogenesis by blocking VEGF-C function. Tissue-specific loss of esKDR in mice induced, at birth, spontaneous lymphatic invasion of the normally alymphatic cornea and hyperplasia of skin lymphatics without affecting blood vasculature. Administration of esKDR inhibited lymphangiogenesis but not hemangiogenesis induced by corneal suture injury or transplantation, enhanced corneal allograft survival and suppressed lymphangioma cellular proliferation. Naturally occurring esKDR thus acts as a molecular uncoupler of blood and lymphatic vessels; modulation of esKDR might have therapeutic effects in treating lymphatic vascular malformations, transplantation rejection and, potentially, tumor lymphangiogenesis and lymphedema.

References

1. Shibata et al, BMC Medicine 8 (2010)
2. Albuquerque et al, Nature Med 2009
3. Ebos et al, Mol Cancer Res 2 (2004)
4. Ebos et al, Cancer res 68 (2008).

Database References Antigen

Protein RefSeq:	NP_002241.1
Uniprot ID:	P35968
mRNA RefSeq:	NM_002253.2

Product Specifications

Species reactivity	human
Clone/Ab feature	IgG ₁ ; #EIC
Cross reactivity	ND
Host	mouse
Clonality	monoclonal
Purification	Protein G purified
Immunogen	recombinant human soluble KDR (D7) (RT# S01-002)
Formulation	liquid
Buffer/Stabilizer	PBS; 1% BSA
Preservative	0,02% sodium azide
Conjugation	FITC

Warnings: Reagents contain sodium azide. Under acidic conditions sodium azide yields hydrazoic acid, this is extremely toxic. Azide compounds should be diluted with running water before discarding. These precautions are recommended to avoid deposits in plumbing where explosive condition may develop.

Storage: Store protected from light at 2-8°C. **Do not freeze!**



DO NOT FREEZE!

Specificity: The unconjugated antibody will detect native human VEGFR-2/KDR in Western analysis, ELISA experiments and on the surface of different human cell types.

Application:

FACS analysis: Use at 2-10 µg/ml.

NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!