



20150120ML

Anti-Human CCL-11 (#11F7)

**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

| | |
|------------------|----------------------------|
| Cat.-no.: | 101-M256 |
| Size: | 100 µg |
| Lot. No.: | According to product label |

Preparation: This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a mouse) immunized with human recombinant protein of CCL-11 (also called Eotaxin).

Target Background

| | |
|---------------------------|---------------|
| Synonyms (Target): | CCL11; SCYA11 |
|---------------------------|---------------|

CCL11 is a potent eosinophil chemoattractant that was originally purified from bronchoalveolar lavage fluid of guinea pigs sensitized by aerosol challenge with ovalbumin. Microsequencing of the purified protein revealed the guinea pig CCL11 to be a member of the beta (CC) chemokine family of inflammatory and immunoregulatory cytokines. cDNA clones for guinea pig, mouse, and human CCL11 have been isolated. Human CCL11 cDNA encodes a 97 amino acid residue precursor protein from which the aminoterminal 23 amino acid residues are cleaved to generate the 74 amino acid residue mature human CCL11. At the protein sequence level, mature human CCL11 is approximately 60% identical to mature mouse and guinea pig CCL11. In addition, human CCL11 also shows high amino acid sequence identity to human MCP1, 2, and 3. Human CCL11 is chemotactic for eosinophils, but not mononuclear cells or neutrophils. The CC chemokine receptor 3 (CCR3) has now been identified to be a specific human CCL11 receptor. CCR3 has also been shown to serve as a cofactor for a restricted subset of primary HIV viruses and binding of CCL11 to CCR3 inhibited infection by the HIV isolates.

Database References Target

| | |
|------------------------|-------------|
| Protein RefSeq: | NP_002977.1 |
| Uniprot ID: | P51671 |
| mRNA RefSeq: | NM_002986.2 |

Product Specifications

| | |
|------------------------------|--------------------------|
| Host | Mouse |
| Reactivity against | Human |
| Clonality | Monoclonal Antibody |
| Clone | (#11F7) |
| Isotype | IgG1 |
| Purification | Protein G chromatography |
| Antigen | recombinant human CCL-11 |
| Formulation | lyophilized |
| Reconstitution buffer | PBS (sterile) |

Reconstitution: Reconstitute the antibody with 200 µl sterile PBS and the final concentration is 500 µg/ml.

Stability: Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C. Reconstituted antibody can be aliquoted and stored frozen at < -20 °C for at least for six months without detectable loss of activity.

Remarks: This antibody was selected for its ability to detect human CCL-11 protein.

**AVOID REPEATED FREEZE AND THAW CYCLES!**

Applications

The antibody can be used within the following applications:

WB, IHC, N

Recommended usage:

Western Blot: 1:500 - 1:1000

Neutralization: Yes

IHC (Paraffin): Yes

NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!