



20150126ML

# Anti-Human CD50 (#8V23)

**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

<b>Cat.-no.:</b>	<b>101-M309</b>
Size:	100 µg
Lot. No.:	According to product label

**Preparation:** This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a mouse) immunized with human recombinant protein of CD50 extracellular domain (also called ICAM-3).

## Target Background

<b>Synonyms (Target):</b>	ICAM3; CD50; CDW50; ICAM-R
---------------------------	----------------------------

Intercellular adhesion molecule 3 (ICAM-3), also known as CD50, is a member of the immunoglobulin superfamily, binds the leukocyte integrins LFA-1 (CD11a/CD18) and alphaD/beta2. ICAM-3 is expressed on leukocytes and epidermal Langerhans cells.

## Database References Target

<b>Protein RefSeq:</b>	NP_002153.2
<b>Uniprot ID:</b>	P32942
<b>mRNA RefSeq:</b>	NM_002162.3

## Product Specifications

<b>Host</b>	Mouse
<b>Reactivity against</b>	Human
<b>Clonality</b>	Monoclonal Antibody
<b>Clone</b>	(#8V23)
<b>Isotype</b>	IgG1
<b>Purification</b>	Protein G chromatography
<b>Antigen</b>	Human recombinant CD50 EC domain
<b>Formulation</b>	lyophilized
<b>Reconstitution buffer</b>	PBS (sterile)

**Reconstitution:** Reconstitute the antibody with 200 µl sterile PBS and the final concentration is 500 µg/ml.

**Stability:** Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C. Reconstituted antibody can be aliquoted and stored frozen at < -20 °C for at least for six months without detectable loss of activity.

**Remarks:** This antibody was selected for its ability to detect human CD50.

**AVOID REPEATED FREEZE AND THAW CYCLES!**

## Applications

The antibody can be used within the following applications:

WB, N

**Recommended usage:**

Western Blot: 1:500 - 1:2000

Neutralization of CD50 bioactivity: Yes

**NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!**