



20190325DS

Anti-Human PDGFR-beta (#5C34)

**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

Cat.-no.:	101-M43A
Size:	100 µg
Lot. No.:	According to product label

Preparation: This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a mouse immunized with recombinant human platelet-derived growth factor receptor alpha extracellular domain).

Target Background

Synonyms (Target):	PDGFRB; JTK12; PDGFR; CD140B; PDGFR1; PDGFR-1
---------------------------	---

PDGF is a major serum mitogen that can exist as a homo or heterodimeric protein consisting of disulfide-linked PDGF-A and PDGF-B chains. The PDGF-AA, PDGF-BB and PDGF-AB isoforms have been shown to bind to two distinct cell surface PDGF receptors with different affinities. Where as PDGF R α binds all three PDGF isoforms with high affinity, PDGF R β binds PDGF-BB only with high affinity. Both PDGF R α and PDGF R β are members of the class III subfamily of receptor tyrosine kinases (RTK) that also includes the receptors for M-CSF, SCF and Flt3 ligand. All class III RTKs are characterized by the presence of five immunoglobulin-like domains in their extracellular region and a split kinase domain in their intracellular region. PDGF binding induces receptor homo- and heterodimerization and signal transduction. The expression of the α and β receptors is independently regulated in various cell types. Recombinant soluble PDGF R β binds PDGF with high affinity and is potent PDGF antagonist.

Database References Target

Protein RefSeq:	NP_002600
Uniprot ID:	P09619
mRNA RefSeq:	NM_002609.3

Product Specifications

Host	Mouse
Reactivity against	Human
Clonality	Monoclonal Antibody
Clone	(#5C34)
Isotype	IgG1
Purification	Protein G chromatography
Antigen	recombinant human PDGFR beta extracellular domain
Formulation	lyophilized
Reconstitution buffer	PBS (sterile)

Reconstitution: Reconstitute the antibody with 200 µl sterile PBS and the final concentration is 500 µg/ml.

Stability: Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C. Reconstituted antibody can be aliquoted and stored frozen at <-20°C for at least for six months without detectable loss of activity.

Remarks: This antibody detects human PDGFR- β in Western blots. No cross-reactivity is shown with human PDGFR- α .

**AVOID REPEATED FREEZE AND THAW CYCLES!**

Applications

The antibody can be used within the following applications:

WB

Recommended usage:**WB:** 1:400-1000

NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!