



Anti-Human IL-17 (CTLA-8) (#9J17)

20230310DS



FOR RESEARCH ONLY! NOT FOR HUMAN USE!

Cat.-no.:	101-M783
Size:	100 µg
Lot. No.:	According to product label

Preparation: This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a mouse) immunized with human recombinant protein of IL-17.

Target Background

Synonyms (Target):	IL17A; IL17; CTLA8; IL-17; IL-17A
---------------------------	-----------------------------------

The IL-17 family is comprised of at least six proinflammatory cytokines that share a conserved cysteine-knot structure but diverge at the N-terminus. IL-17 family members are glycoproteins secreted as dimers that induce local cytokine production and recruit granulocytes to sites of inflammation. IL-17 is induced by IL-15 and IL-23, mainly in activated CD4+ T cells distinct from Th1 or Th2 cells. IL-17F is the most homologous to IL-17, but is induced only by IL-23 in activated monocytes. IL-17B binds the IL-17B receptor, but not the IL-17 receptor; it is most homologous with IL-17D, which is expressed by resting CD4+ T cells and CD19+ B cells. IL-17E is mainly produced by Th2 cells and recruits eosinophils to lung tissue. IL-17C has a very restricted expression pattern but has been detected in adult prostate and fetal kidney libraries.

Database References Target

Protein RefSeq:	NP_002181.1
Uniprot ID:	Q16552
mRNA RefSeq:	NM_002190.2

Product Specifications

Host	Mouse
Reactivity against	Human
Clonality	Monoclonal Antibody
Clone	(#9J17)
Isotype	IgG1
Purification	Protein G chromatography
Antigen	Human recombinant CTLA-8 (IL-17)
Formulation	lyophilized
Reconstitution buffer	PBS

Application/Handling

Reconstitution: Centrifuge vial prior to opening. Reconstitute the antibody with 500 µl sterile PBS and the final concentration is 200 µg/ml.

Stability: Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C. Reconstituted antibody can be aliquoted and stored frozen at < -20°C for at least for six months without detectable loss of activity.



AVOID REPEATED FREEZE AND THAW CYCLES!

Applications

The antibody can be used within the following applications:

IHC, N, Capture Ab

NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!