



20150116ML

Anti-Mouse IGF-2 (#3J42)

**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

Cat.-no.:	103-M241
Size:	100 µg
Lot. No.:	According to product label

Preparation: This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a rat immunized with purified mouse recombinant protein of IGF-2. The IgG2 fraction of the culture supernatant was purified by Protein A/G affinity chromatography.

Target Background

Synonyms (Target):	Igf2; Mpr; M6pr; Peg2; Igf-2; Igf-II; AL033362
---------------------------	--

Insulin-like growth factor (IGF)-I (also known as somatomedin C and somatomedin A) and IGF-II (multiplication stimulating activity or MSA) belong to the family of insulin-like growth factors that are structurally homologous to proinsulin. Mature IGF-I and IGF-II share approximately 70% sequence identity. Both IGF-I and IGF-II are expressed in many tissues and cell types and may have autocrine, paracrine and endocrine functions. Mature IGF-I and IGF-II are highly conserved between the human, bovine and porcine proteins (100% identity), and exhibit cross-species activity.

Database References Target

Protein RefSeq:	NP_001116208.1
Uniprot ID:	P90535
mRNA RefSeq:	NM_001122736.1

Product Specifications

Host	Rat
Reactivity against	Mouse
Clonality	Monoclonal Antibody
Clone	(#3J42)
Isotype	IgG2
Purification	Protein G/A chromatography
Antigen	recombinant mouse protein of IGF-II
Formulation	lyophilized
Reconstitution buffer	PBS (sterile)

Reconstitution: Reconstitute the antibody with 200 µl sterile PBS and the final concentration is 500 µg/ml.

Stability: Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C. Reconstituted antibody can be aliquoted and stored frozen at < -20 °C for at least for six months without detectable loss of activity.

Remarks: This antibody detects mouse IGF-II in Western blotting. It is cross-reactive with human IGF-II as well.

**AVOID REPEATED FREEZE AND THAW CYCLES!**

Applications

The antibody can be used within the following applications:

WB

Recommended usage:

WB: 1:500-1000

NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!