



# Anti-Human IL-17E

20150223ML



**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

<b>Cat.-no.:</b>	<b>102-P163</b>
Size:	100 µg
Lot. No.:	According to product label

**Preparation:** Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-17E (human IL-17E). Anti-hIL-17E specific antibody was purified by affinity chromatography employing immobilized hIL-17E matrix.

## Target Background

<b>Synonyms (Target):</b>	IL25; IL17E
---------------------------	-------------

The IL-17 family is comprised of at least six proinflammatory cytokines that share a conserved cysteine-knot structure but diverge at the N-terminus. IL-17 family members are glycoproteins secreted as dimers that induce local cytokine production and recruit granulocytes to sites of inflammation. IL-17 is induced by IL-15 and IL-23, mainly in activated CD4+ T cells distinct from Th1 or Th2 cells. IL-17F is the most homologous to IL-17, but is induced only by IL-23 in activated monocytes. IL-17B binds the IL-17B receptor, but not the IL-17 receptor; it is most homologous with IL-17D, which is expressed by resting CD4+ T cells and CD19+ B cells. IL-17E is mainly produced by Th2 cells and recruits eosinophils to lung tissue. IL-17C has a very restricted expression pattern but has been detected in adult prostate and fetal kidney libraries.

### Database References Target

<b>Protein RefSeq:</b>	NP_073626.1
<b>Uniprot ID:</b>	Q9H293
<b>mRNA RefSeq:</b>	NM_022789

## Product Specifications

<b>Species reactivity</b>	Human
<b>Clone/Ab feature</b>	Rabbit IgG
<b>Cross reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal Antibody
<b>Purification</b>	Antigen-affinity purified
<b>Immunogen</b>	Recombinant human IL-17E
<b>Formulation</b>	lyophilized from PBS
<b>Reconstitution buffer</b>	water

**Reconstitution:** Reconstitute the antibody in sterile water to a concentration of 0.1 - 1.0 mg/ml.

**Stability:** The lyophilized antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C.



**AVOID REPEATED FREEZE AND THAW CYCLES!**

### Applications

**Western Blot:** To detect hIL-17E by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-17E is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

**ELISA:** To detect h hIL-17E by direct ELISA (using 100 µl/well antibody solution) a concentration of at least 0.5 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hIL-17E.

**NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!**