

RELIATech GmbH
Lindenerstr. 15
38300 Wolfenbüttel
Germany

Tel.: +49 5331 8586 987
Fax.: +49 5331 8586 989
Email: info@reliatech.de
web: www.reliatech.de

Antigen-Affinity Purified Polyclonal Antibodies

Anti-murine Eotaxin-2

Description: Produced from sera of goats pre-immunized with highly pure (>98%) recombinant murine Eotaxin-2 (murine Eotaxin-2). Anti-murine Eotaxin-2 specific antibody was purified by affinity chromatography employing immobilized murine Eotaxin-2 matrix.

Host species:	Goat
Antigen:	Recombinant murine Eotaxin-2
Purification:	Affinity chromatography
Stabilizer:	none
Buffer:	0.5 x PBS; pH 7.4
Formulation:	lyophilized

Reconstitution: Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Stability: The lyophilized antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. **Avoid repeated freeze-thaw cycles.**

Applications

Western Blot: To detect murine Eotaxin-2 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant murine Eotaxin-2 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Neutralization: To yield one-half maximal inhibition [ND₅₀] of the biological activity of mEotaxin-2 (100 ng/ml), a concentration of 7.0-8.0 µg/ml of this antibody is required.

ELISA: Indirect: To detect murine Eotaxin-2 by indirect ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant murine Eotaxin-2.

Sandwich: To detect murine Eotaxin-2 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant murine Eotaxin-2.

Usage: Anti murine Eotaxin-2 is offered for research use. Not for drug use. **Not for human use.**

Catalogue number: 103-P48G

Size: 100 µg

****Please note: always centrifuge product before opening vial.****