



# Anti-Mouse IL-17A

20150223ML



**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

<b>Cat.-no.:</b>	<b>103-P76</b>
Size:	100 µg
Lot. No.:	According to product label

**Preparation:** Produced from sera of goats pre-immunized with highly pure (98%) recombinant murine IL-17A. Anti-murine IL-17A specific antibody was purified by affinity chromatography employing immobilized murine IL-17A matrix.

## Target Background

<b>Synonyms (Target):</b>	Il17a; Il17; Ctla8; IL-17; Ctla-8; IL-17A
---------------------------	---

The antibody reacts with mouse and rat IL-17A with no recognition of IL-17F. Interleukin-17A (IL-17A) is a CD4+ T cell-derived cytokine that promotes inflammatory responses in cell lines and is elevated in rheumatoid arthritis, asthma, multiple sclerosis, psoriasis, and transplant rejection. The cDNA encoding human IL-17A was isolated from a library of CD4+ T cells; the encoded protein exhibits 72 percent amino acid identity with HVS13, an open reading frame from a T lymphotropic Herpesvirus saimiri, and 63 percent with mouse CTLA-8 (cytotoxic T-lymphocyte associated antigen-8). Human IL-17A exists as glycosylated 20-30 kD homodimers. High levels of IL-17A homodimer are produced by activated peripheral blood CD4+ T-cells. IL-17A enhances expression of the intracellular adhesion molecule-1 (ICAM-1) in human fibroblasts. Human IL-17A also stimulates epithelial, endothelial, or fibroblastic cells to secrete IL-6, IL-8, G-CSF, and PGE2. In the presence of human IL-17A, fibroblasts can sustain the proliferation of CD34+ hematopoietic progenitors and induce maturation into neutrophils. Mouse, rat, and human IL-17A can induce IL-6 secretion in mouse stromal cells, indicating that all homologs can recognize the mouse IL-17A receptor.

### Database References Target

<b>Protein RefSeq:</b>	NP_034682.1
<b>Uniprot ID:</b>	Q62386
<b>mRNA RefSeq:</b>	NM_010552.3

## Product Specifications

<b>Species reactivity</b>	Mouse
<b>Clone/Ab feature</b>	Rabbit IgG
<b>Cross reactivity</b>	Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal Antibody
<b>Purification</b>	Antigen-affinity purified
<b>Immunogen</b>	Recombinant mouse IL-17A
<b>Formulation</b>	lyophilized from PBS
<b>Reconstitution buffer</b>	water

**Reconstitution:** Reconstitute the antibody in sterile water to a concentration of 0.1 - 1.0 mg/ml.

**Stability:** The lyophilized antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C.



**AVOID REPEATED FREEZE AND THAW CYCLES!**

### Applications

**Western Blot:** To detect IL-17A by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIL-15 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

#### ELISA:

**Indirect:** To detect IL-17A by indirect ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant IL-17A.

**Sandwich:** To detect IL-17A by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with an appropriate secondary conjugated antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant IL-17A.

**NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!**