



20150116ML

Anti-Mouse IL-15 (#15B9)

**FOR RESEARCH ONLY! NOT FOR HUMAN USE!**

Cat.-no.:	103-M409
Size:	100 µg
Lot. No.:	According to product label

Preparation: This antibody was produced from a hybridoma (mouse myeloma fused with spleen cells from a rat) immunized with mouse recombinant protein of IL-15.

Target Background

Synonyms (Target):	IL15; IL-15; AI503618
---------------------------	-----------------------

IL-15 is an immunomodulating cytokine that stimulates the proliferation of T lymphocytes and shares many biological properties with IL-2. IL-15 exerts its biological activities primarily on T cells. It is also essential in the development, survival and activation of NK cells. Increased expression of IL-15 has been implicated with rheumatoid arthritis, inflammatory bowel disease and diseases affiliated with retroviruses HIV and HTLV-I. Human IL-15 is biologically active on mouse cells as measured by the dose-dependent stimulation of the proliferation of mouse CTLL cells. Recombinant murine IL-15 is a 13.3 kDa protein consisting of 115 amino acid residues.

Database References Target

Protein RefSeq:	NP_032383.1
Uniprot ID:	P48346
mRNA RefSeq:	NM_008357

Product Specifications

Host	Rat
Reactivity against	Mouse
Clonality	Monoclonal Antibody
Clone	(#15B9)
Isotype	IgG2
Purification	Protein A/G chromatography
Antigen	recombinant mouse protein of IL-15
Formulation	lyophilized
Reconstitution buffer	PBS (sterile)

Reconstitution: Reconstitute the antibody with 200 µl sterile PBS and the final concentration is 500 µg/ml.

Stability: Lyophilized samples are stable for 2 years from date of receipt when stored at -70°C. Reconstituted antibody can be aliquoted and stored frozen at < -20 °C for at least for six months without detectable loss of activity.

Remarks: Remarks: This antibody detects specifically mouse IL-15 with WB.

**AVOID REPEATED FREEZE AND THAW CYCLES!**

Applications

The antibody can be used within the following applications:
WB

Recommended usage:

WB: 1:50-200

NOTE: OPTIMAL DILUTIONS SHOULD BE DETERMINED BY EACH LABORATORY FOR EACH APPLICATION!